

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

THIS PAGE BLANK (USPTO)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) EP 0 752 472 A1

(12)

**EUROPEAN PATENT APPLICATION**

published in accordance with Art. 158(3) EPC

(43) Date of publication:

08.01.1997 Bulletin 1997/02

(51) Int. Cl.<sup>6</sup>: C12N 15/31, C12N 1/21,  
C12P 13/08, C12P 13/14

(21) Application number: 95909970.6

(86) International application number:  
PCT/JP95/00269

(22) Date of filing: 23.02.1995

(87) International publication number:  
WO 95/23224 (31.08.1995 Gazette 1995/37)(84) Designated Contracting States:  
CH DE ES FR GB IT LI NL

(30) Priority: 24.02.1994 JP 26501/94

(71) Applicant: Ajinomoto Co., Inc.  
Tokyo 104 (JP)

(72) Inventors:

- KIMURA, Eiichiro  
Ajinomoto Co., Inc.  
Technology &  
Kawasaki-ku, Kawasaki-shi Kanagawa 210 (JP)
- ABE, Chizu  
Ajinomoto Co., Inc.  
Technology &  
Kawasaki-ku, Kawasaki-shi Kanagawa 210 (JP)

- KAWAHARA, Yoshio  
Ajinomoto Co., Inc.  
Technology &  
Kawasaki-ku, Kawasaki-shi Kanagawa 210 (JP)
- YOSHIHARA, Yasuhiko  
Ajinomoto Co., Inc.  
Technology  
Kawasaki-ku, Kawasaki-shi Kanagawa 210 (JP)
- NAKAMATSU, Tsuyoshi  
Ajinomoto Co., Inc.  
Technology  
Kawasaki-ku, Kawasaki-shi Kanagawa 210 (JP)

(74) Representative: Kolb, Helga, Dr. Dipl.-Chem. et al  
Hoffmann, Eitle & Partner,  
Patent- und Rechtsanwälte,  
Arabellastrasse 4  
81925 München (DE)**(54) NOVEL GENE ORIGINATING IN CORYNEBACTERIUM AND USE THEREOF**

(57) The L-lysine productivity of an L-lysine-producing corynebacterium is enhanced by amplifying a novel gene originating in a corynebacterium and participating in L-glutamic acid production, while the L-glutamic acid productivity of an L-glutamic acid-producing corynebacterium is enhanced by suppressing the function of the above gene.




Fig. 1

**Description**Field of The Invention

5 The present invention relates to breeding and use of Coryneform bacteria which are used for the fermentative production of L-amino acids such as L-glutamic acid and L-lysine and other substances, and the use thereof.

Description of the Prior Art

10 When Coryneform bacteria are cultivated in a medium having a limited amount of biotin, they produce a large amount of L-glutamic acid. On the other hand, when the Coryneform bacteria are cultivated in a medium containing an excess amount of biotin, they do not produce L-glutamic acid. However, it is known that if a surfactant or penicillin is added to a medium containing such an excess amount of biotin, the growth of the bacteria in this medium is inhibited and the bacteria produce a large amount of L-glutamic acid therein.

15 To make Coryneform bacteria produce L-glutamic acid, any of the following means is effective.

1. The biotin concentration in the medium is made suboptimal. Refer to S. Okumura, T. Tsugawa, T. Tsunoda and A. Kitai, Nippon Nogeikagaku Kaishi, 36, 197-203 (1962).

2. A surfactant is added to the medium provided that a sufficient amount of biotin is present. Refer to I. Shiio, H. Otsuka and N. Atsuya, J. Biochem., 53, 333-340 (1963); K. Takinami, H. Okada and T. Tsunoda, Agr. Biol. Chem., 27, 853-863 (1963).

3. Penicillin is added to the medium provided that a sufficient amount of biotin is present. Refer to U.S. Patent 3,080,297; Japanese Patent Publication No. 37-1695 (1962); M. Shibui, T. Kurima, S. Okabe and T. Osawa, Amino Acid and Nucleic Acid, 17, 61-65 (1968).

25 The mechanisms in these means have been considered in the following way.

With respect to L-glutamic acid production by limitation of biotin, the major factor is considered as follows: Biotin is a coenzyme for acetyl-CoA carboxylase for the synthesis of fatty acids and, in addition, unsaturated fatty acids, such as oleic acid, and their derivatives have a substitutive effect for biotin. Therefore, biotin will have an influence on the composition of fatty acids constituting the cell membrane, thereby varying the permeability of L-glutamic acid through the cell membrane (I. Shiio, S. Otsuka and M. Takahashi, J. Biochem., 51, 56-62 (1962); I. Shiio, K. Narui, N. Yahaba and M. Takahashi, J. Biochem., 51, 109-111 (1962)).

The production of L-glutamic acid by addition of a surfactant or penicillin has also been considered with reference to the variation in the permeability of L-glutamic acid through the cytoplasmic membrane because of the variation in the structure of cell surface (Shiio, S. Otsuka and N. Katsuya, J. Biochem., 53, 333-340 (1963)).

As mentioned above, the production of L-glutamic acid has been discussed with reference to its permeability through cell membrane, but no finding directly verifying the relationship therebetween has been obtained.

There have been many unclear matters with regard to by what mechanisms the limitation of biotin or the addition of a surfactant or penicillin improves the ability of Coryneform bacteria to produce L-glutamic acid.

40 In addition, no information on the gene level, which will be an important key factor for clarifying these mechanisms, has been available.

Disclosure of the Invention

45 The object of the present invention is to clarify the mechanism of L-glutamic acid-production possessed by Coryneform bacteria, specifically the function of the surfactant to be added in the mechanism of L-glutamic acid production possessed by Coryneform bacteria, and to breed and improve Coryneform bacteria producing L-glutamic acid and the like on the basis of the information thus obtained.

Specifically, the object of the present invention is to clarify, on the gene level, the mechanisms of L-glutamic acid-production by Coryneform bacteria, to isolate a gene of Coryneform bacteria that participates in surfactant resistance, and to apply the thus-obtained gene to the breeding of L-glutamic acid-producing Coryneform bacteria and to the production of L-glutamic acid and the like by Coryneform bacteria.

50 The present inventors have intensively studied so as to attain the above-mentioned object and, as a result, have found the existence of a gene which will participate in the L-glutamic acid-production by Coryneform bacteria (the gene is hereinafter referred to as dtSR gene, and the protein encoded by the gene is referred to as DTSR protein). In addition, we have found a valuable use for this gene. On the basis of these findings, we have completed the present invention.

The present invention includes the following embodiments:

(1) A gene derived from a Coryneform bacterium and coding for a protein which imparts surfactant resistance to

said bacterium.

(2) The gene according to (1), wherein the protein comprises an amino acid sequence of amino acid number 37 to 543 in the amino acid sequence shown by SEQ ID NO: 2 in the Sequence Listing, or an amino acid sequence having, in the amino acid sequence, substitution, deletion or insertion which does not substantially adversely affect on activity to impart the surfactant resistance to Coryneform bacteria.

(3) The gene according to (1), which has a sequence of from 467th to 1987th nucleotides in the nucleotide sequence shown by SEQ ID NO: 1 in the Sequence Listing, or a nucleotide sequence which is substantially the same as the sequence.

(4) A recombinant DNA obtainable by ligating a vector which functions in Coryneform bacteria to the gene as defined in (1), (2) or (3).

(5) A Coryneform bacterium harboring the recombinant DNA as defined in (4).

(6) A method for producing L-lysine comprising cultivating a Coryneform bacterium harboring the recombinant DNA as defined in (4) and capable of producing L-lysine in a liquid medium to produce and accumulate L-lysine in the culture, and collecting the L-lysine.

(7) A gene comprising a nucleotide sequence having substitution, deletion, insertion, addition or inversion of one or more nucleotides in a nucleotide sequence of the gene as defined in (1), (2) or (3) so that a protein encoded by the nucleotide sequence does not normally function regarding an activity to impart surfactant resistance to Coryneform bacteria.

(8) A recombinant DNA obtainable by ligating a vector which functions in Coryneform bacteria to the gene as defined in (7).

(9) A Coryneform bacterium having substitution, deletion, insertion, addition or inversion of one or more nucleotides in a nucleotide sequence of the gene as defined in (1), (2) or (3) or a promoter thereof on chromosome so that a protein having an activity to impart surfactant resistance to Coryneform bacteria does not normally function.

(10) A method for producing L-glutamic acid comprising cultivating a Coryneform bacterium having substitution, deletion, insertion, addition or inversion of one or more nucleotides in a nucleotide sequence of the gene as defined in (1), (2) or (3) or a promoter thereof on chromosome so that a protein having an activity to impart surfactant resistance to Coryneform bacteria does not normally function, and capable of producing L-glutamic acid, in a liquid medium to produce and accumulate L-glutamic acid in the culture, and collecting the L-glutamic acid.

The present invention will be described in detail hereinafter.

Coryneform bacteria as referred to herein are a group of microorganisms defined in Bergey's Manual of Determinative Bacteriology, 8th Ed., p. 599 (1974). The bacteria are aerobic, Gram-positive, non-acid-fast bacilli not having the ability to sporulate, and include bacteria which had been classified as bacteria belonging to the genus Brevibacterium but have now been unified into the genus Corynebacterium [see Int. J. Syst. Bacteriol., 41, 255 (1981)] and also include bacteria of the genus Brevibacterium and Microbacterium which are closely related to the genus Corynebacterium. Of such Coryneform bacteria, those mentioned below, which are known as L-glutamic acid-producing bacteria, are most preferred for use in the present invention.

Corynebacterium acetoacidophilum

Corynebacterium acetoglutamicum

Corynebacterium callunae

Corynebacterium glutamicum

Corynebacterium lilium (Corynebacterium glutamicum)

Corynebacterium melassecola

Brevibacterium divaricatum (Corynebacterium glutamicum)

Brevibacterium lactofermentum (Corynebacterium glutamicum)

Brevibacterium saccharolyticum

Brevibacterium immariophilum

Brevibacterium roseum

Brevibacterium flavum (Corynebacterium glutamicum)

Brevibacterium thiogenitalis

Specifically, the following strains of these bacteria are exemplified:

<u>Corynebacterium acetoacidophilum</u>	ATCC 13870
<u>Corynebacterium acetoglutamicum</u>	ATCC 15806
<u>Corynebacterium callunae</u>	ATCC 15991
<u>Corynebacterium glutamicum</u>	ATCC 13032
<u>Corynebacterium glutamicum</u>	ATCC 13060

<u>Brevibacterium divaricatum</u>	ATCC 14020
<u>Brevibacterium lactofermentum</u>	ATCC 13869
<u>Corynebacterium lilium</u>	ATCC 15990
<u>Corynebacterium melassecola</u>	ATCC 17965
5 <u>Brevibacterium saccharolyticum</u>	ATCC 14066
<u>Brevibacterium immariophilum</u>	ATCC 14068
<u>Brevibacterium roseum</u>	ATCC 13825
<u>Brevibacterium flavum</u>	ATCC 13826
10 <u>Brevibacterium thioogenitalis</u>	ATCC 19240

These strains can be obtained from the American Type Culture Collection (ATCC). A registration number has been assigned to each strain of bacteria. Based on the registration number, anyone can obtain the corresponding strain of bacteria from ATCC. The registration numbers of the strains of bacteria deposited in ATCC are described in the ATCC catalog.

15 Surfactants as referred in the present invention function to accelerate the production of L-glutamic acid by Coryneform bacteria, like penicillin, in a medium containing an excess amount of biotin therein, and these include various non-ionic surfactants, cationic surfactants and anionic surfactants [see K. Yamada, J. Takahashi and J. Nakamura, the Hakkokogaku Kaishi, 20, 348-350 (1962); K. Udagawa, S. Abe and I. Kinoshita, Hakkokogaku Kaishi, 40, 614-619 (1962)]. Of nonionic surfactants, Tween 60 (polyoxyethylene sorbitan monostearate) and Tween 40 (polyoxyethylene sorbitan monopalmitate) have a penicillin-like effect [see I. Shiio, S. Otsuka and N. Katsuya, J. Biochem., 53, 333-340 (1963)]. C<sub>3</sub> to C<sub>18</sub> free saturated fatty acids have the same effect by themselves [see K. Takinami, H. Okada and T. Tsunoda, Agr. Biol. Chem., 28, 114-118 (1964)]. Tween 40 was used in the examples of the present invention.

#### (1) Isolation of a gene from a Coryneform bacterium participating in surfactant resistance

25 To isolate a gene from a Coryneform bacterium participating in surfactant resistance, for example, the following process can be employed.

- 30 (1) A surfactant-sensitive mutant of a Coryneform bacterium which shows higher sensitivity with regard to surfactants is obtained;
- (2) various fragments of a chromosomal DNA prepared from a wild type strain of a Coryneform bacterium are each ligated to a vector that functions in Coryneform bacteria to produce various recombinant DNAs;
- (3) the recombinant DNAs each are introduced into cells of the surfactant-sensitive mutant of a Coryneform bacterium to conduct transformation;
- 35 (4) from the resulting transformants, strains which have lost the surfactant-sensitivity are selected;
- (5) the recombinant DNAs are recovered from the thus-selected surfactant-insensitive transformants; and
- (6) the structure of the chromosomal DNA fragment of the wild type strain of a Coryneform bacterium ligated to the vector is analyzed.

40 The chromosomal DNA fragment of the wild type strain of a Coryneform bacterium thus obtained contains a gene derived from a Coryneform bacterium participating in surfactant resistance. This gene participates at least in the mechanism of the accumulation of L-glutamic acid by Coryneform bacteria in a medium containing a surfactant. In addition, this gene also participates in the production of L-glutamic acid in a medium containing penicillin or containing a limited amount of biotin.

45 A surfactant-sensitive mutant of a Coryneform bacterium which shows higher sensitivity to surfactants means a mutant of a Coryneform bacterium which grows poorly in a medium containing a surfactant at such a concentration that does not have any influence on the growth of the wild type strain of Coryneform bacteria in the medium. Regarding a surfactant of polyoxyethylene sorbitan monopalmitate, a surfactant-sensitive mutant of a Coryneform bacterium grows worse than the corresponding wild type strain in a medium containing the surfactant at a concentration of from 0.1 to 1 mg/dl. On the contrary, the growth of a wild type strain of a Coryneform bacterium is not affected by the presence of the surfactant at a concentration of from 0.1 to 1 mg/dl in the medium. When such a mutant is cultivated in a medium containing an excess amount of biotin to produce L-glutamic acid therein by adding a surfactant, the necessary concentration of the surfactant to be added to the medium may be lower than that in the ordinary case. It is considered that the condition of the cells of the surfactant-sensitive mutant will be similar to that of the cells of the corresponding wild type strain which are exposed to the surfactants.

55 To obtain a surfactant-sensitive mutant of a Coryneform bacterium, the method described in Japanese Patent Application Laid-Open No. 50-126877 (1975) (Japanese Patent Publication No. 52-24593 (1977)) can be employed.

As one example of the surfactant-sensitive mutant of a Coryneform bacterium, mentioned is *Brevibacterium lactofermentum* (*Corynebacterium glutamicum*) AJ 11060. This mutant was deposited in the National Institute of Bioscience

and Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry, under the accession number FERM P-3678.

To prepare various fragments of the chromosomal DNA of a wild type strain of a Coryneform bacterium, the following process may be employed. The wild type strain of a Coryneform bacterium is cultivated in a liquid medium, the cells grown therein are collected, and the chromosomal DNA is recovered from the collected cells according to the method of Saito et al. [see H. Saito and K. Miura, *Biochem. Biophys. Acta* **72**, 619 (1963)]. The thus-recovered chromosomal DNA is partially cleaved with restriction enzymes. Four-base recognition enzymes are used as the restriction enzymes, and the cleavage is conducted to prepare various DNA fragments under the condition under which the DNA is incompletely decomposed.

The vector functioning in Coryneform bacteria as referred to herein is, for example, a plasmid which is autonomously replicable in Coryneform bacteria. Specific examples of the vector are mentioned below.

pAM 330	see Japanese Patent Application Laid-Open No. 58-67699 (1983)
pHM 1519	see Japanese Patent Application Laid-Open No. 58-77895 (1983)
pAJ 655	see Japanese Patent Application Laid-Open No. 58-192900 (1983)
pAJ 611	see Japanese Patent Application Laid-Open No. 58-192900 (1983)
pAJ 1844	see Japanese Patent Application Laid-Open No. 58-192900 (1983)
pCG 1	see Japanese Patent Application Laid-Open No. 57-134500 (1982)
pCG 2	see Japanese Patent Application Laid-Open No. 58-35197 (1983)
pCG 4	see Japanese Patent Application Laid-Open No. 57-183799 (1982)
pCG 11	see Japanese Patent Application Laid-Open No. 57-183799 (1982)

To prepare various recombinant DNAs by ligating the vector functioning in Coryneform bacteria and various fragments of the chromosomal DNA of a wild type strain of Coryneform bacteria, the following process may be employed. The vector is first cleaved with a restriction enzyme. The restriction enzyme to be used for cleaving the vector is the same as that used for cleaving the chromosomal DNA, or is such one by which the vector is cleaved to give ends complementary to the ends of the fragment of the chromosomal DNA. The ligation of the vector and the DNA fragment is generally effected via a ligase, such as T4 DNA ligase, etc.

To introduce the recombinant DNA to the surfactant-sensitive mutant of a Coryneform bacterium, any known transformation methods can be employed. For instance, employable are a method of treating recipient cells with calcium chloride so as to increase the permeability of DNA, which has been reported for *Escherichia coli* K-12 [see Mandel, M. and Higa, A., *J. Mol. Biol.*, **53**, 159 (1970)]; and a method of preparing competent cells from cells which are at the growth phase followed by introducing the DNA thereinto, which has been reported for *Bacillus subtilis* [see Duncan, C.H., Wilson, G.A. and Young, F.E., *Gene*, **1**, 153 (1977)]. In addition to these, also employable is a method of making DNA-recipient cells into the protoplast or spheroplast which can easily take up recombinant DNAs followed by introducing the recombinant DNA into the cells, which is known to be applicable to *Bacillus subtilis*, actinomycetes and yeasts [see Chang, S. and Choen, S.N., *Molec. Gen. Genet.*, **168**, 111 (1979); Bibb, M.J., Ward, J.M. and Hopwood, O.A., *Nature*, **274**, 398 (1978); Hinnen, A., Hicks, J.B. and Fink, G.R., *Proc. Natl. Sci., USA*, **75**, 1929 (1978)].

The above-mentioned protoplast method for *Bacillus subtilis* can be employed in the present invention to obtain sufficiently high-level frequency. In addition to this, however, also employable is a method of introducing a DNA into protoplast of cells of a Coryneform bacterium while the protoplast are kept in contact with either polyethylene glycol or polyvinyl alcohol and with divalent metal ions, as described in Japanese Patent Application Laid-Open No. 57-183799 (1982). In this method, carboxymethyl cellulose, dextran, Ficoll, Pluronic F68 (produced by Serva Co.), etc. may also be used, in place of polyethylene glycol or polyvinyl alcohol, so as to accelerate the introduction of DNA to the protoplast cells. In the examples of the present invention, the transformation was conducted by an electric pulse method (see Japanese Patent Application Laid-Open No. 2-207791 (1990)).

The method for selecting the strain which has lost the surfactant sensitivity from the transformants will be described below.

DNA fragments having a size of approximately from 4 to 6 kbp, which have been obtained by partially digesting the chromosomal DNA of a wild type strain of a Coryneform bacterium with the restriction enzyme, *Sau3AI*, are ligated to a plasmid vector which is autonomously replicable both in *Escherichia coli* and in Coryneform bacteria to construct recombinant DNAs, and these recombinant DNAs are introduced into the competent cells of *Escherichia coli* DH5 (produced by Takara Shuzo Co., Ltd.). The resulting transformants are cultivated to prepare a gene library of the wild type strain of a Coryneform bacterium.

By using the recombinant DNA in this gene library, *Brevibacterium lactofermentum* AJ 11060 is transformed. The resulting transformants are inoculated on a surfactant-free M-CM2G agar plate (containing 5 g of glucose, 10 g of polypeptone, 10 g of yeast extract, 5 g of NaCl, 0.2 g of DL-methionine, 15 g of agar and 4 mg of chloramphenicol in one liter of water, and having a pH of 7.2), and about 40,000 colonies are formed thereon. These colonies are replicated onto a M-CM2G plate containing 30 mg/liter of a surfactant, Tween 40, and the colonies growing well on this surfactant-

containing M-CM2G plate are collected. Thus, transformants which have lost the surfactant sensitivity are obtained.

To recover the recombinant DNAs from the thus-obtained transformants which have lost the surfactant sensitivity, the same method as that employed for preparing the chromosomal DNA of the wild type strain of a Coryneform bacterium may be employed. Briefly, the transformants are cultivated in a liquid medium, the cells are collected from the culture, and the recombinant DNAs are recovered from them according to the method of Saito et al. [see H. Saito and K. Miura, *Biochem. Biophys. Acta* **72**, 619 (1963)].

The structure of the chromosomal DNA fragment of the wild type strain of a Coryneform bacterium that has been ligated to the vector is analyzed as follows. The full-length nucleotide sequence of the chromosomal DNA fragment is determined by the dideoxy method which is an ordinary nucleotide sequencing method, and the structure of the DNA is analyzed to determine the positions of the enhancer, the promoter, the operator, the SD sequence, the leader peptide, the attenuator, the initiation codon, the termination codon, the open reading frame, etc.

The gene obtained from the Coryneform bacterium participating in surfactant resistance is dtsR gene, which has a sequence ranging between the 467th to 469th nucleotides (ATG) and 1985th to 1987th nucleotides (CTG) shown by SEQ ID NO: 1 in the Sequence Listing. The amino acid sequence which may be encoded by the gene is shown in SEQ NO: 1 and SEQ NO: 2 in the Sequence Listing. Another ATG (nucleotide number 359-361) is present at the upstream of the ATG of 467th to 469th nucleotides in the same frame and possibility of this ATG being an initiation codon is not denied. However, from analysis of consensus sequence in the upstream region of the gene, it is presumed that the ATG of 467th to 469th nucleotides is an initiation codon. That is, the amino acid sequence of amino acid number 37 to 543 in the amino acid sequence shown by SEQ ID NO: 2 is presumed to be the amino acid sequence of the DTSR protein. When the amino acid sequence of the DTSR protein and the nucleotide sequence of dtsR gene are referred to in the present specification and claims, these may be described with ATG of 467th to 469th nucleotides as the initiation codon. However, it should be considered possibility of the ATG of 359th to 361st nucleotides being the initiation codon. For example, if the dtsR gene is introduced to a Coryneform bacterium to enhance the expression, expression of the sequence of nucleotide number 467 to 1987 in the nucleotide sequence shown by SEQ ID NO: 1 is considered to be sufficient. However, one skilled in the art will easily appreciate that when the coding region and the upstream region of the nucleotide sequence shown by SEQ ID NO: 1 including nucleotide number 359-466 are introduced to a Coryneform bacterium, the DTSR protein can be properly expressed whichever ATG is the initiation codon. In either case, the N-terminal methionine coded by the initiation codon can be cleaved with an aminopeptidase in the expression of the dtsR gene in the cells.

According to the search on data base, it has been confirmed that the dtsR gene having the nucleotide sequence shown by SEQ ID NO: 1 and the DTSR protein encoded by this sequence are novel. It has been found that this protein is homologous to the protein described in *Proc. Natl. Acad. Sci. USA*, **83**, 8049-8053 (1986); *Proc. Natl. Acad. Sci. USA*, **83**, 4864-4869 (1986); and *Gene*, **122**, 199-202 (1992), as propionyl-CoA carboxylase (PCC)  $\beta$  subunit protein. However, none of these references suggests that the protein participates in the production of glutamic acid.

Propionyl-CoA carboxylase is an enzyme that catalyzes one reaction in the metabolic pathway in which  $\alpha$ -ketoglutaric acid is converted into succinyl-CoA via 2-hydroxyglutaric acid, propionyl-CoA, D-methylmalonyl-CoA and L-methylmalonyl-CoA, and it seems that the metabolic pathway is a by-pass route for the reaction to be catalyzed by  $\alpha$ -ketoglutarate dehydrogenase in the TCA cycle. In this connection, it should be specifically noted that propionyl-CoA carboxylase is an enzyme needing biotin as a coenzyme, indicating that the production of glutamic acid in the surfactant-addition method is related to the production of glutamic acid in the biotin-limitation method.

## (2) Preparation of a Coryneform bacterium having the recombinant DNA

The recombinant DNA containing the dtsR gene from a Coryneform bacterium participating in surfactant resistance, which is obtained in the foregoing item (1) is prepared in vitro, and is introduced into Coryneform bacteria. By the introduction, a Coryneform bacterium can be prepared in which the intracellular concentration of the DTSR protein is increased. The general means for this purpose is to enhance the intracellular expression of the dtsR gene or to increase the number of copies of the dtsR gene in the cells.

To enhance the intracellular expression of the dtsR gene, the gene is ligated downstream of a strong promoter. The dtsR gene is exemplified by a nucleotide sequence which codes for an amino acid sequence of amino acid number 37 to 543 in the amino acid sequence shown by SEQ ID NO: 2. Specifically, a nucleotide sequence of nucleotide number 467 to 1987 in the nucleotide sequence shown by SEQ ID NO: 1 and a nucleotide sequence which is substantially the same as the nucleotide sequence are mentioned. The term "nucleotide sequence which is substantially the same" used herein means a nucleotide sequence coding for a protein which is substantially the same as the protein encoded by the nucleotide sequence of nucleotide number 467 to 1987 with respect to the activity to impart surfactant resistance to Coryneform bacteria. As to these nucleotide sequences, the coded DTSR protein may have substitution, deletion or insertion of an amino acid which does not substantially affect on the activity to impart surfactant resistance to Coryneform bacteria.

As strong promoters functioning intracellularly in Coryneform bacteria, known are the lac promoter, tac promoter



and Trp promoter derived from *Escherichia coli* [see Y. Morinaga, M. Tsuchiya, K. Miwa and K. Sano, J. Biotech., 5, 305-312 (1987)]. The trp promoter derived from a Coryneform bacterium is also preferred (see Japanese Patent Application Laid-Open No. 62-195294 (1987)).

The DNA containing the *dtbR* gene and the DNA containing such a promoter are prepared separately, and these are ligated to each other in vitro. To cleave these DNAs and to ligate them together, restriction enzymes and a ligase are employed, respectively. The recombinant DNA obtained by the ligation is then introduced into cells of a Coryneform bacterium. For the introduction, the same process as that referred to in the foregoing item (1) can be employed.

To introduce the recombinant DNA comprising a strong promoter and the *dtbR* gene into cells of a Coryneform bacterium, a vector functioning in the cells of the Coryneform bacterium must be used. When the vector referred to in item (1) is used in this step, the recombinant DNA is held outside the chromosome. In order to make the recombinant DNA hold onto the chromosomal DNA, a temperature-sensitive plasmid such as that described in Japanese Patent Application Laid-Open No. 5-7491 (1993) is used as the vector, and the cells are cultivated at a nonpermissible temperature to cause homologous recombination.

The expression of the *dtbR* gene by the thus-obtained cells of the Coryneform bacterium having therein the recombinant DNA comprising a strong promoter and the *dtbR* gene is enhanced and the intracellular concentration of the DTSR protein is increased.

When the DNA containing the *dtbR* gene is ligated to a multi-copy plasmid and introduced into cells of a Coryneform bacterium, the number of copies of the said gene in the cells is increased. As examples of such a multi-copy plasmid, those mentioned in item (1) are referred to.

Also, employable is a process for causing homologous recombination by using, as a target, a sequence that exists in large numbers on the chromosomal DNA of a Coryneform bacterium. As one example of the sequence much existing on the chromosomal DNA of the Coryneform bacterium, mentioned is an insertion sequence existing at both ends of the transposal element of the Coryneform bacterium. The sequence and a method of causing such homologous recombination by using the sequence are disclosed in International Publication No. WO 93/18151.

The expression of the *dtbR* gene by the thus-obtained cells of a Coryneform bacterium in which the number of copies of the *dtbR* gene has been increased is enhanced, and the intracellular concentration of the DTSR protein is increased.

### 3) Production of L-lysine by Coryneform bacteria having the recombinant DNA therein

Various artificial mutants have heretofore been used as L-lysine-producing bacteria. Using these as the hosts, the recombinant DNA of the present invention can be introduced into them to improve their L-lysine productivity. Such artificial mutants are as follows: S-(2-aminoethyl)-cysteine (hereinafter referred to as "AEC")-resistant mutants; mutants requiring amino acids such as L-homoserine for their growth (see Japanese Patent Publication Nos. 48-28078 (1973) and 56-6499 (1981)); mutants resistant to AEC and requiring amino acids such as L-leucine, L-homoserine, L-proline, L-serine, L-arginine, L-alanine, L-valine, etc. (see U.S. Patent Nos. 3,708,395 and 3,825,472); L-lysine-producing mutants resistant to DL- $\alpha$ -amino- $\epsilon$ -caprolactam,  $\alpha$ -amino-lauryl-lactam, aspartic acid analogues, sulfa drugs, quinoids and N-lauroyl-leucine, and L-lysine-producing mutants resistant to oxaloacetate decarboxylase inhibitors or respiratory system enzyme inhibitors (see Japanese Patent Application Laid-Open Nos. 50-53588 (1975), 50-31093 (1975), 52-102498 (1977), 53-9394 (1978), 53-86089 (1978), 55-9783 (1980), 55-9759 (1980), 56-32995 (1981), 56-39778 (1981), Japanese Patent Publication Nos. 53-43591 (1978) and 53-1833 (1978)); L-lysine-producing mutants requiring inositol or acetic acid (see Japanese Patent Application Laid-Open Nos. 55-9784 (1980) and 56-8692 (1981)); L-lysine-producing mutants sensitive to fluoropyruvic acid or to temperatures of 34°C or higher (see Japanese Patent Application Laid-Open No. 53-86090 (1978)); L-lysine-producing mutants of *Brevibacterium* or *Corynebacterium* resistant to ethylene glycol (see U.S. Patent No. 4,411,997).

As specific examples, the following strains are referred to.

*Brevibacterium lactofermentum* AJ 12031 (FERM BP-277); see Japanese Patent Application Laid-Open No. 60-62994 (1985).

*Brevibacterium lactofermentum* ATCC 39134; see Japanese Patent Application Laid-Open No. 60-62994 (1985).

*Corynebacterium glutamicum* AJ 3463 (FERM P-1987); see Japanese Patent Publication No. 51-34477 (1976).

*Brevibacterium lactofermentum* AJ 12435 (FERM BP-2294); see U.S. Patent No. 5,304,476.

*Brevibacterium lactofermentum* AJ 12592 (FERM BP-3239); see U.S. Patent No. 5,304,476.

*Corynebacterium glutamicum* AJ 12596 (FERM BP-3242); see U.S. Patent No. 5,304,476.

In the Coryneform bacterium obtained by introducing the recombinant DNA of the present invention into these L-lysine-producing bacteria according to the process in the foregoing item 2) the intracellular concentration of the DTSR protein is increased, and the resultant bacterium has the ability to produce a large amount of L-lysine.

The medium used for the production of L-lysine may be ordinary medium containing carbon sources, nitrogen

sources, inorganic ions and, if desired, other minor organic nutrients.

Saccharides such as hydrolysates of starch, etc., alcohols such as ethanol, inositol, etc.; organic acids such as acetic acid, fumaric acid, citric acid, succinic acid, etc are usable.

As the nitrogen sources, usable are inorganic ammonium salts such as ammonium sulfate, ammonium chloride, ammonium phosphate, etc.; organic nitrogen compounds such as hydrolysates of soybeans, etc.; as well as ammonia gas, aqueous ammonia, etc.

As the inorganic ions, small amounts of potassium phosphate, magnesium sulfate, iron ions, manganese ions, etc. are added to the medium. As the minor organic nutrients, essential substances such as vitamin B<sub>1</sub>, etc., as well as yeast extract, etc are preferably contained in suitable amounts, if necessary.

It is recommended that the cultivation is performed under an aerobic condition for 16 to 72 hours at a temperature ranging from 30 to 45°C, and during the cultivation, the pH of the media is controlled to between 5 and 7. To adjust the pH, inorganic or organic, acidic or alkaline substances, ammonia gas, etc. can be used.

The collection of L-lysine from the fermentation liquid may be conducted by an ordinary ion-exchange method, a precipitation method and other known methods in combination.

#### 4) Preparation of Coryneform bacteria in which the DTSR protein does not normally function

The *dtsR* gene has been obtained as a gene that imparts resistance to surfactants to a Coryneform bacteria, as mentioned in the foregoing item (1). Therefore, it was expected that the strain with the amplified *dtsR* gene would no longer produce L-glutamic acid in a medium to which a surfactant was added at a concentration at which the wild type strain of a Coryneform bacterium produces L-glutamic acid in the presence of an excess amount of biotin. In consideration of this, the effect of the amplification of the *dtsR* gene in the production of L-glutamic acid by adding a surfactant was investigated according to the method shown in the examples described below, and, as was expected, the noticeable inhibition of the production of L-glutamic acid was observed. In addition, it was confirmed that the amplification of the *dtsR* gene also resulted in the suppression of the production of L-glutamic acid in the biotin-limiting method and in the penicillin-adding method in the presence of an excess amount of biotin. These results indicate that the *dtsR* gene does not only make the strain resistant to surfactants but also plays an important role in the production of L-glutamic acid.

For these reasons, it was expected that if, opposite to the above, the expression of the *dtsR* gene is inhibited and the intracellular concentration or activity of the DTSR protein is lowered, the ability of the cells to produce L-glutamic acid might be improved and L-glutamic acid might be produced especially in a medium containing an excess amount of biotin without adding any biotin activity-suppressing substance such as surfactants or antibiotics. In consideration of this, a strain in which the *dtsR* gene of the wild type strain was disrupted was prepared and the ability of the thus-modified strain to produce L-glutamic acid was examined, and it has been confirmed that the *dtsR* gene-disrupted mutant strain produced a large amount of L-glutamic acid even if biotin is present at a high concentration at which the wild type strain produces little L-glutamic acid, as is demonstrated in the examples as described below.

The strain in which the *dtsR* gene is mutated can be obtained by a method inducing mutation by use of a chemical agent or by a breeding method using recombinant DNA technique. When the gene has already been obtained, the recombinant DNA technique is employed, whereby the gene can easily be disrupted by homologous recombination. The means of disrupting a gene by homologous recombination has already been established. For this, employable are a method of using a linear DNA and a method of using a temperature-sensitive plasmid.

Specifically, by site-specific mutation [see Kramer, W. and Faits, H.J., *Methods in Enzymology*, 154, 350 (1987)] or mutation with chemicals such as sodium hyposulfite, hydroxylamine, etc. [see Shortle, D. and Nathans, D., *Proc. Natl. Acad. Sci. USA*, 75, 270 (1978)], the substitution, deletion, insertion, addition or inversion of one or more nucleotides is made in the nucleotide sequence of the coding region or of the promoter region in the *dtsR* gene. The thus-modified or disrupted gene is substituted for the normal gene on the chromosome, thereby lowering or vanishing the activity of the genetic product, DTSR protein, or lowering or vanishing the transcription of the gene.

In the site-specific mutagenesis, synthetic oligonucleotides are used, and according to this method, it is possible to introduce any desired substitution, deletion, insertion, addition or inversion to only (a) limited base pair(s). To carry out this method, a plasmid having the intended gene which has been cloned and whose nucleotide sequence has been determined, is first denatured to prepare a single-stranded DNA. Next, a synthetic oligonucleotide complementary to the site to be mutated is prepared. The synthetic oligonucleotide shall be such that it does not have a completely complementary sequence but may have any desired nucleotide substitution, deletion, insertion, addition or inversion. After this, the single-stranded DNA is annealed with the synthetic oligonucleotide having such a desired nucleotide substitution, deletion, insertion, addition or inversion. Then, this is formed into a complete double-stranded plasmid, using the Klenow fragment of DNA polymerase I and T4 ligase. The resulting plasmid is then introduced into the competent cell of *Escherichia coli*. Some of these transformants thus obtained have a plasmid containing the gene that has the desired nucleotide substitution, deletion, insertion, addition or inversion fixed therein. As a similar method for modifying or disrupting the gene by introducing the mutation, known is the recombinant PCR method [see PCR Technology, Stockton

Press (1989)].

The chemical treatment method is a method for directly treating the DNA fragment containing the intended gene with sodium hyposulfite, hydroxylamine or the like chemical thereby randomly introducing the mutation having nucleotide substitution, deletion, insertion, addition or inversion into the DNA fragment.

Whether or not the desired mutation is introduced can be confirmed by transforming a surfactant-sensitive mutant with the DNA fragment subjected to mutation treatment and determining the presence of surfactant resistance of the resultant transformant. On this occasion, by examining growth both at high temperature and at low temperature within temperature at which the Coryneform bacteria can usually grow, and selecting a transformant which can grow at the low temperature and the growth of which is inhibited at higher temperature, a mutated gene which becomes temperature-sensitive can be obtained.

As the means of substituting the thus-obtained gene that has been modified or disrupted by the introduction of the mutation for the normal gene on the chromosome of a Coryneform bacterium, employable is a method of using homologous recombination [see Experiments in Molecular Genetics, Cold Spring Harbor Laboratory Press (1972); Matsuyama, S. and Mizushima, S., J. Bacteriol., 162, 1196 (1985)]. According to homologous recombination, when a plasmid or the like having a sequence which is homologous to the sequence of the chromosome is introduced into the cell, a recombination occurs at the sites having the homologous sequence at a certain frequency whereby the complete plasmid introduced is incorporated into the chromosome. After this, when a further recombination occurs at the sites having the homologous sequence on the chromosome, the plasmid is removed from the chromosome. By the latter recombination, the gene into which the mutation has been introduced is fixed onto the chromosome, depending on the recombined site, and the original normal gene may be removed from the chromosome along with the plasmid. By selecting such a strain, it is possible to obtain a strain in which the gene modified or disrupted by the introduction thereof of the mutation having nucleotide substitution, deletion, insertion, addition or inversion has been substituted for the normal gene on the chromosome.

#### 5) Production of L-glutamic acid by Coryneform bacteria in which the DTSR protein does not normally function

As described above, a strain which is the L-glutamic acid-producing Coryneform bacterium and in which intracellular concentration or activity the DTSR protein is lowered, that is, a strain in which the DTSR protein does not normally function is improved in productivity of L-glutamic acid. Especially, the strain can produce L-glutamic acid without addition of biotin activity-suppressing substance such as a surfactant or antibiotic even if biotin is present in an excess amount.

For obtaining a mutant in which the DTSR protein does not normally function by breeding Coryneform bacteria having the ability to produce L-glutamic acid, a glutamic acid-producing wild type strain or a mutant derived therefrom of Coryneform bacteria can be used as the starting strain. As examples of the mutant, the following are mentioned:

Brevibacterium lactofermentum AJ 12745 (FERM-BP 2922); see U.S. Patent No. 5,272,067.

Brevibacterium lactofermentum AJ 12746 (FERM-BP 2923); see U.S. Patent No. 5,272,067.

Brevibacterium flavum AJ 12747 (FERM-BP 2924); see U.S. Patent No. 5,272,067.

Corynebacterium glutamicum AJ 12478 (FERM-BP 2925); see U.S. Patent No. 5,272,067.

Corynebacterium glutamicum ATCC 21492.

The medium to be used for the production of L-glutamic acid may be ordinary medium containing carbon sources, nitrogen sources, inorganic ions and, if desired, other minor organic nutrients.

As the carbon sources, usable are saccharides such as glucose, lactose, galactose, fructose, hydrolysates of starch, etc.; alcohols such as ethanol, inositol, etc.; organic acids such as acetic acid, fumaric acid, citric acid, succinic acid, etc.

As the nitrogen sources, usable are inorganic ammonium salts such as ammonium sulfate, ammonium chloride, ammonium phosphate, etc.; organic nitrogen compounds such as hydrolysates of soybeans, etc.; as well as ammonia gas, aqueous ammonia, etc.

As the inorganic ions, small amounts of potassium phosphate, magnesium sulfate, iron ions, manganese ions, etc. are added to the medium. As the minor organic nutrients, essential substances such as vitamin B<sub>1</sub>, etc., as well as yeast extract, etc. are preferably contained in suitable amounts, if necessary.

It is recommended that the cultivation is carried out under an aerobic condition for 16 to 72 hours at a temperature ranging from 30 to 45°C, and during the cultivation, the pH of the medium is kept at from 5 to 8. To adjust the pH, inorganic or organic, acidic or alkaline substances, ammonia gas, etc. can be used.

Surfactants or penicillin may be added to the medium where the *dtsR* gene-disrupted strain thus obtained is cultivated, or the biotin concentration in the medium may be limited. In this way, the yield of L-glutamic acid to be produced in the medium may be increased further.

The collection of L-glutamic acid from the fermentation liquid may be effected by an ordinary ion exchange method,

precipitation method and other known methods in combination.

#### Brief Explanation of the Drawings

5 Fig. 1 shows the growth of AJ 11060/pDTR6 (*dtsR* gene-amplified strain) and AJ 11060/pSAC4 (control) in a surfactant-free or surfactant-added medium.

- : pDTR6-introduced strain in surfactant-free medium
- : pSAC4-introduced strain in surfactant-free medium
- 10 ○ : pDTR6-introduced strain in surfactant-added medium
- : pSAC4-introduced strain in surfactant-added medium

Fig. 2 shows the degree of surfactant resistance of ATCC 13869/pHSGX-KΔE (deletion-mutation type *dtsR* gene-amplified strain), ATCC 13869/pDTR6 (*dtsR* gene-amplified strain) and ATCC 13869/pSAC4 (control).

- 15 ○ : pSAC4-introduced strain
- △ : pHSGX-KΔE-introduced strain
- △ : pDTR6-introduced strain

20 Fig. 3 shows the growth curve of AJ 11060/pDTR6 and AJ 11060/pSAC4 in a medium containing 300 or 3 μg/liter of biotin.

- : pDTR6-introduced strain in the presence of 300 μg/liter of biotin
- : pSAC4-introduced strain in the presence of 300 μg/liter of biotin
- 25 ○ : pDTR6-introduced strain in the presence of 3 μg/liter of biotin
- : pSAC4-introduced strain in the presence of 3 μg/liter of biotin

#### Description of the Preferred Embodiment

30 The present invention will be described in more detail with reference to the following examples.

#### Example 1 (Preparation of chromosomal DNA of *Brevibacterium lactofermentum* ATCC 13869 (a wild type strain of *Coryneform* bacteria))

35 *Brevibacterium lactofermentum* ATCC 13869 was inoculated in 100 ml of a T-Y medium [containing 1% Bacto-trip-ton (by Difco), 0.5% Bacto-yeast extract (by Difco) and 0.5% NaCl; pH 7.2] and cultivated at 31.5°C for 8 hours to obtain a culture. This culture was subjected to centrifugation at 3,000 r.p.m. for 15 minutes to obtain 0.5 g of wet cells. From the wet cells, obtained was the chromosomal DNA according to Saito & Miura method (see Biochem. Biophys. Acta. 72, 619, (1963)). Next, 60 μg of the chromosomal DNA and 3 units of a restriction enzyme, *Sau*3AI were each mixed in 10 mM Tris-HCl buffer (containing 50 mM NaCl, 10 mM MgSO<sub>4</sub> and 1 mM dithiothreitol; pH 7.4), and the reaction was carried out at 37°C for 30 minutes. After the reaction, the reaction mixture was subjected to ordinary phenol extraction and ethanol precipitation to obtain 50 μg of chromosomal DNA fragments of *Brevibacterium lactofermentum* ATCC 13869 digested with *Sau*3AI.

#### Example 2 (Preparation of gene library of *Brevibacterium lactofermentum* ATCC 13869, using plasmid vector DNA)

20 μg of a plasmid vector DNA (pSAC4) capable of autonomously replicating in both the cells of *Escherichia coli* and the cells of *Coryneform* bacteria and 200 units of a restriction enzyme, *Bam*HI were mixed in 50 mM Tris-HCl buffer (containing 100 mM NaCl and 10 mM magnesium sulfate; pH 7.4) and reacted at 37°C for 2 hours to obtain a digested solution, and the solution was then subjected to ordinary phenol extraction and ethanol precipitation. After this, the DNA fragment was dephosphorylated with bacterial alkaline phosphatase according to the method described in Molecular Cloning, 2nd Edition [J. Sambrook, E.F. Fritsch and T. Maniatis, Cold Spring Harbor Laboratory Press, p.1.56 (1989)] so as to prevent the re-binding of the plasmid vector-derived DNA fragment, and then the thus-dephosphorylated DNA fragment was subjected to ordinary phenol extraction and ethanol precipitation.

55 One μg of this pSAC4 digested with *Bam*HI, 1 μg of the chromosomal DNA fragments of *Brevibacterium lactofermentum* ATCC 13869 digested with *Sau*3AI, that had been obtained in Example 1, and 2 units of T4 DNA ligase (produced by Takara Shuzo Co., Ltd.) were added to 66 mM Tris-HCl buffer (pH 7.5) containing 66 mM magnesium chloride, 10 mM dithiothreitol and 10 mM ATP and reacted therein at 16°C for 16 hours to conduct the ligation of the DNA. Next, cells of *Escherichia coli* DH5 were transformed with said DNA mixture by an ordinary method, and the resulting trans-

formant cells were inoculated onto an L-agar medium containing 170 µg/ml of chloramphenicol to obtain about 20,000 colonies constituting a gene library.

### Example 3 (Transformation of *Brevibacterium lactofermentum* AJ 11060)

From these approximately 20,000 colonies mentioned above, the recombinant DNAs were recovered according to the above-mentioned Saito and Miura method.

The recombinant DNA mixture which was divided into 50 batches, was introduced into cells of the strain AJ 11060, a mutant being sensitive to surfactants, by ordinary transformation using electric pulse (see Japanese Patent Application Laid-Open No. 2-207791 (1990)). The resulting transformant cells were inoculated onto a glucose-added L-agar medium and the cultivation was performed by static incubation at 31.5°C, whereby about 20,000 colonies of transformants were formed on the medium. Next, these transformant colonies were replicated to the same plate medium but containing 30 mg/liter of the surfactant, and several strains that were resistant to the surfactant and grown on the plate medium were obtained therefrom.

### Example 4 (Measurement of surfactant resistance of strains having multi-copies of *dtsR* gene)

The recombinant DNA was extracted from each of the several strains that had been grown, and *Brevibacterium lactofermentum* AJ 11060 was re-transformed with the DNA. From the resulting transformants, one surfactant-resistant strain was selected. The recombinant DNA of this strain is designated as pDTR6, and the gene which is carried by this plasmid and which has the ability to make the strain resistant to the surfactant is designated as *dtsR*. The inhibition of the growth of AJ 11060, into which the plasmid had been introduced, in a surfactant(3 g/liter)-added liquid medium is suppressed (see Fig. 1).

### Example 5 (Preparation of DNA)

The plasmid was prepared from AJ 11060/pDTR6 having the recombinant DNA that had been obtained in the above, by an ordinary method, and this was introduced into *Escherichia coli* JM 109. The resulting *Escherichia coli* JM 109/pDTR6 was cultivated in 20 ml of a medium containing 1% of tryptone, 0.5% of yeast extract and 0.5% of NaCl, at 37°C for 24 hours, and 20 ml of the resulting culture was inoculated in one liter of a medium having the same composition as above and cultivation was performed at 37°C for 3 hours. Then, 0.2 g of chloramphenicol was added thereto, and the cultivation was continued for additional 20 hours at the same temperature to obtain a culture. Next, the resulting culture was centrifuged at 3,000 r.p.m. for 10 minutes to obtain 2 g of wet cells. The obtained cells were suspended in 20 ml of 350 mM Tris-HCl buffer (pH 8.0) containing 25% of sucrose, and then 10 mg of lysozyme (produced by Sigma Co.), 8 ml of 0.25 M EDTA solution (pH 8.0) and 8 ml of 20% sodium dodecylsulfate solution were added thereto. Then, the resulting suspension was heated at 60°C for 30 minutes to obtain a lysate. 13 ml of 5 M NaCl solution was added to the lysate, and the lysate was then treated at 4°C for 16 hours. After the treatment, this was centrifuged at 15,000 r.p.m. for 30 minutes. The supernatant thus obtained was subjected to ordinary phenol extraction and ethanol precipitation to obtain a DNA precipitate.

The precipitate was dried under reduced pressure and then dissolved in 6 ml of 10 mM Tris-HCl buffer (pH 7.5) containing 1 mM EDTA, and 6 g of cesium chloride and 0.2 ml of ethidium bromide (19 mg/ml) were added thereto. Then, this was subjected to equilibrium density gradient centrifugation, using an ultracentrifugator, at 39,000 r.p.m. for 42 hours, by which the DNA was isolated. Next, ethidium bromide was removed from this, using n-butanol, and thereafter this was subjected to dialysis against 10 mM Tris-HCl (pH 7.5) containing 1 mM EDTA to obtain about 500 µg of a purified recombinant DNA, pDTR6. A private number AJ 12967 was assigned to *Escherichia coli* JM109/pDTR6. This strain was deposited on February 22, 1994 in National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry, with the accession number FERM P-14168, and transferred on February 9, 1995 to the international deposit under the Budapest Treaty with the accession number FERM BP-4994.

### Example 6 (Analysis of nucleotide sequence of DNA having *dtsR* gene)

The nucleotide sequence of the recombinant DNA obtained in Example 5 was determined. The sequencing was effected according to the Sanger method, using a Taq DyeDeoxy Terminator Cycle Sequencing Kit (produced by Applied Biochemical Co.). The nucleotide sequence of the DNA thus obtained is shown by SEQ ID NO: 1 in the Sequence Listing. While the longest open reading frame in this sequence is a nucleotide sequence of from 359th A to 1987th G in the nucleotide sequence shown by SEQ ID NO: 1, it is presumed from analysis of consensus sequence in the upstream region of the gene that ATG of 467th to 469th is an initiation codon. The amino acid sequence which can be encoded by the open reading frame of 359th A to 1987th G is shown by SEQ ID NO: 1 in the Sequence Listing

together with the nucleotide sequence. Further, only the amino acid sequence is shown by SEQ ID NO: 2 in the Sequence Listing. A protein encoded by the nucleotide sequence of from 467th to 1987th nucleotides is designated as the DTSR protein.

It is well known that the N-terminal methionine residue of protein is deleted after translation by the action of peptidase. This is because the N-terminal methionine is derived from the translation initiation codon, ATG, and therefore has no relation to the intrinsic function of protein in most cases. Also in the DTSR protein of the present invention, there is a probability that the methionine residue is deleted.

The nucleotide sequence and the amino acid sequence were compared with known sequences with respect to the homology. The data bases used for the comparison were EMBL and SWISS-PROT. As a result, it has been confirmed that the gene represented by SEQ ID NO: 1 in the Sequence Listing and the protein encoded by this are novel.

#### Example 7 (Confirmation of participation of *dtsR* gene in surfactant resistance)

The open reading frame was identified by the nucleotide sequencing, which suggested the existence of DTSR protein. In order to further confirm that the gene which imparts surfactant resistance to Coryneform bacteria exists in this region, a part of gene in the region of the open reading frame was deleted in an in-frame manner from the gene fragment represented by SEQ ID NO: 1 in the Sequence Listing, and this gene fragment was investigated as to whether or not it has an activity of imparting surfactant resistance to Coryneform bacteria. Specifically, pDTR6 was digested with *Xba*I and *Kpn*I to obtain a fragment containing *dtsR* gene. This gene fragment was ligated to a fragment of plasmid pHSG398 (produced by Takara Shuzo Co., Ltd.) that had been treated with *Xba*I and *Kpn*I, using T4 DNA ligase (produced by Takara Shuzo Co., Ltd.), to prepare plasmid pHSGX-K. The *dtsR* gene has two sites that are digested with *Eco*52I, at the nucleotides 766 and 1366 in SEQ ID NO: 1. pHSGX-K was completely digested with *Eco*52I and then self-ligated to prepare plasmid pHSGX-KΔE from which 600 base pairs of the *Eco*52I fragments were deleted. The structure of the *dtsR* gene on this pHSGX-KΔE was such that the center part was deleted in an in-frame manner.

Next, in order to make pHSGX-KΔE capable of autonomously replicating in Coryneform bacteria, the replication origin derived from a known plasmid, pHM1519, capable of self-amplification in Coryneform bacteria (see Miwa, K. et al., Agric. Biol. Chem. 48, 2901-2903 (1984); Japanese Patent Application Laid-Open No. 5-7491 1993)) was introduced into the only one *Kpn*I cleavage site existing on pHSGX-KΔE. Specifically, pHM1519 was digested with restriction enzymes, *Bam*H1 and *Kpn*I to obtain a gene fragment having a replication origin, and the resulting fragment was made to have blunt ends, using a blunting kit produced by Takara Shuzo Co., Ltd, and then introduced into the *Kpn*I site of pHSGX-KΔE, using a *Kpn*I linker (produced by Takara Shuzo Co., Ltd.), to obtain pKCX-KΔE. As a control sample, the replication origin of pHM1519 was inserted into the *Sal*I site of pHSG399, using *Sal*I linker (produced by Takara Shuzo Co., Ltd.) to prepare pSAC4. These pKCX-KΔE and pSAC4 prepared herein were separately introduced into cells of a wild type strain of Coryneform bacteria, *Brevibacterium lactofermentum* ATCC 13869, according to the above-mentioned electric pulse method, and the resulting transformants were examined with respect to the degree of surfactant resistance. For the examination, the cells were cultivated in a M-CM2G liquid medium, to which from 0 to 10 mg/dl of polyoxyethylene sorbitan monopalmitate had been added, and the degree of the cell growth in the culture was measured.

As a result, it was verified that the deletion-type *dtsR* gene had lost the function to impart the surfactant resistances.

#### Example 8 (Production of L-glutamic acid using *dtsR* gene-amplified strain by biotin-limiting method)

The strain AJ 11060/pDTR6 was cultivated to produce L-glutamic acid, according to the biotin-limiting method, as mentioned below. Specifically, the strain AJ 11060/pDTR6 was separately cultivated on an M-CM2G plate medium containing 4 mg/liter of chloramphenicol. Then, the grown cells were inoculated in a medium containing 80 g of glucose, 1 g of  $\text{KH}_2\text{PO}_4$ , 0.4 g of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 30 g of  $(\text{NH}_4)_2\text{SO}_4$ , 0.01 g of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.01 g of  $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ , 15 ml of soybean hydrolysate solution, 200  $\mu\text{g}$  of thiamine hydrochloride, 60  $\mu\text{g}$  of biotin, 4 mg of chloramphenicol and 50 g of  $\text{CaCO}_3$  in one liter of pure water (the pH of the medium having been adjusted to 7.0 with KOH) and cultivated therein at 31.5°C for 20 hours. The resulting culture was inoculated in the same medium as above but not containing biotin (this medium is hereinafter referred to as a "biotin-limited medium"), at a concentration of 5% by volume, and the cultivation was carried out at 31.5°C for about 20 hours.

While cultivating the strain AJ 11060/pDTR6 in the biotin-limited medium, the amount of biotin required by the unit weight of the cells grown in the medium was measured (see Fig. 3). As a control, the strain AJ 11060/pSAC4 was cultivated in the same manner as above.

The amounts of the grown cells of the strain AJ 11060/pDTR6 were larger than that of the control. Thus, the biotin demand per unit weight of the grown cells of the AJ 11060/pDTR6 was lowered than that of the strain AJ 11060/pSAC4.

Example 9 (Production of L-lysine by dtsR gene-amplified strain)

pDTR6 was introduced into an L-lysine-producing strain of Coryneform bacteria, *Brevibacterium lactofermentum* AJ 12435 (FERM BP-2294), by an electric pulse method. The resulting transformant was cultivated in a medium mentioned below to produce L-lysine.

Specifically, the strain was cultivated on an M-CM2G plate medium containing 4 mg/liter of chloramphenicol, and the thus-obtained cells were inoculated in a medium containing 100 g of glucose, 1 g of  $\text{KH}_2\text{PO}_4$ , 0.4 g of  $\text{MgSO}_4$ , 30 g of  $(\text{NH}_4)_2\text{SO}_4$ , 0.01 g of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.01 g of  $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ , 15 ml of soybean hydrolysate solution, 200  $\mu\text{g}$  of thiamine hydrochloride, 300  $\mu\text{g}$  of biotin, 4 mg of chloramphenicol and 50 g of  $\text{CaCO}_3$  in one liter of pure water (the pH of the medium having been adjusted to 7.0 with KOH) at 32°C for 40 hours. As a control, a strain into which pSAC4 is introduced in the same manner instead of pDTR6 was used. The results are shown in Table 5.

Table 5

Strain	L-lysine (g/liter)
AJ 12435/pSAC4	15
AJ 12435/pDTR6	25

Reference Example 1 (Production of L-glutamic acid using dtsR gene-amplified strain by surfactant-added method)

By adding a surfactant as a biotin activity-suppressing agent, the strain AJ 11060 can produce a large amount of L-glutamic acid even if a high concentration of biotin is present. However, since *dtsR* gene-amplified strains have elevated surfactant resistance, it was considered that the production of glutamic acid by these strains will be decreased, even though a surfactant is added. The strains AJ 11060/pSAC4 and AJ 11060/pDTR6 were cultivated to produce L-glutamic acid, according to the surfactant-added method mentioned below. Specifically, the strains were each cultivated on an M-CM2G plate medium containing 4 mg/liter of chloramphenicol. Then, the cells were inoculated onto a medium containing 80 g of glucose, 1 g of  $\text{KH}_2\text{PO}_4$ , 0.4 g of  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 30 g of  $(\text{NH}_4)_2\text{SO}_4$ , 0.01 g of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.01 g of  $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ , 15 ml of soybean hydrolysate solution, 200  $\mu\text{g}$  of thiamine hydrochloride, 300  $\mu\text{g}$  of biotin, 4 mg of chloramphenicol, 3.0 g of polyoxyethylene sorbitan monopalmitate and 50 g of  $\text{CaCO}_3$  in one liter of pure water (the pH of the medium having been adjusted to 8.0 with KOH) and cultivated therein at 31.5°C for 20 hours.

After the cultivation, the amount of L-glutamic acid produced and accumulated in each culture was measured. The yields obtained are shown in Table 1.

Table 1

Strain	Yield of L-glutamic Acid (%)
AJ 11060/pSAC4	29.6
AJ 11060/pDTR6	9.0

By amplifying the *dtsR* gene, the yield of L-glutamic acid remarkably reduced. This indicates that the DTSR protein closely involves in the production of L-glutamic acid by the surfactant-added method.

Reference Example 2 (Production of L-glutamic acid using dtsR gene-amplified strain by penicillin-added method)

The strains AJ 11060/pSAC4 and AJ 11060/pDTR6 were cultivated to produce L-glutamic acid by the penicillin-added method in the same manner as in Reference Example 1, except that 30 units of penicillin G was added to the medium in place of polyoxyethylene sorbitan monopalmitate. The results are shown in Table 2.

Table 2

Strain	Yield of L-glutamic Acid (%)
AJ 11060/pSAC4	27.6
AJ 11060/pDTR6	12.8

By amplifying the *dtsR* gene, the yield of L-glutamic acid remarkably reduced. This indicates that the DTSR protein also involves in the production of L-glutamic acid by the penicillin-added method, as well as the production of L-glutamic acid by the surfactant-added method.

#### Example 10 (Production of *dtsR* gene-disrupted strain)

Since it was found that the production of L-glutamic acid was decreased by the amplification of the *dtsR* gene, it was expected that the disruption of the *dtsR* gene would result in an increase in the yield of L-glutamic acid. The gene-disrupted strain was obtained by the homologous recombination method disclosed in Japanese Patent Application Laid-Open No. 5-7491 (1993) using a temperature-sensitive plasmid.

Specifically, the replication origin which had been obtained from a plasmid capable of autonomously replicating in Coryneform bacteria and in which the autonomous replication had become temperature-sensitive, was introduced into the *KpnI* recognition site of pHSGX-KΔE shown in Example 7, to form plasmid pKTCX-KΔE.

This pKTCX-KΔE was introduced into a wild type strain, *Brevibacterium lactofermentum* ATCC 13869 by an electric pulse method, and the *dtsR* gene on the chromosome was substituted by a deleted-type *dtsR* gene according to the method described in Japanese Patent Application Laid-Open No. 5-7491 (1993). Specifically, ATCC 13869/pKTCX-KΔE was cultivated by shaking in a M-CM2G liquid medium containing 50 μg/ml of oleic acid at 25°C for 6 hours, and the culture was inoculated onto an M-CM2G medium containing 5 μg/ml of chloramphenicol and 50 μg/ml of oleic acid. The plasmid-inserted strains formed colonies at 34°C, and these were collected. From the thus-obtained strains, those that had become sensitive to chloramphenicol at 34°C were selected by a replication method. The chromosomes of these sensitive strains were obtained by an ordinary method, and the structure of the *dtsR* gene of each of these chromosomes was analyzed by the Southern hybridization method. Thus, the strain in which the *dtsR* gene had been substituted by a deleted-type *dtsR* gene was confirmed and designated as a strain ΔE.

#### Example 11 (Estimation of L-Glutamic acid productivity of the strain ΔE)

The strains, ATCC 13869, AJ 11060 and ΔE, were cultivated to produce L-glutamic acid in the manner mentioned below. These strains were each refreshed by cultivating on an M-CM2G plate medium. The thus-refreshed strains were inoculated into a medium containing 80 g of glucose, 1 g of KH<sub>2</sub>PO<sub>4</sub>, 0.4 g of MgSO<sub>4</sub>, 30 g of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.01 g of FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.01 g of MnSO<sub>4</sub>·7H<sub>2</sub>O, 15 ml of soybean hydrolysate, 200 μg of thiamine hydrochloride, 300 μg of biotin, 1 g of Tween 80 (polyoxyethylene sorbitan monooleate) and 50 g of CaCO<sub>3</sub> in one liter of pure water (the pH of the medium having been adjusted to 8.0 with KOH), and the cultivation was carried out at 31.5°C for 20 hours. The results are shown in Table 3.

Table 3

Strain	L-Glutamic Acid (g/liter)
ATCC 13869	0
AJ 11060	0
ΔE	34

With respect to the wild type strain and the strain AJ 11060, the accumulation of L-glutamic acid was not observed due to biotin present in an excess amount in the medium. In contrast, the strain ΔE well produced and accumulated L-glutamic acid.



Industrial Applicability

The dtsR gene of the present invention is a gene which plays an important role in the production of L-glutamic acid in Coryneform bacteria which are used for the production of L-glutamic acid by fermentation. When this gene is amplified in L-lysine-producing Coryneform bacteria, the L-lysine productivity is improved. When this gene is disrupted in L-glutamic acid-producing Coryneform bacteria, the L-glutamic acid productivity is improved.

10

15

20

25

30

35

40

45

50

55

## SEQUENCE LISTING

## INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2855

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

## (ii) MOLECULAR TYPE: Genomic DNA

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 359..1987

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

15	GATCTTGGAA	CTCGACAGTT	TTCACCGTCC	AGTTTGGAGC	GCCTGAGCTT	GCAAGCTCCA	60
	GCAAGTCAGC	ATTAGTGGAG	CCTGTCACCT	TTTCGTAAAT	GACCTGGCCA	AAGTCACCGT	120
	TTTGGAGCAA	TTTTTCCTTC	AGGAGCTCAA	CGTTTAGCGG	CTCTCTGGAT	CGTGAAATGT	180
	CAACGTTT	CAT GGAAGCCAAT	GTAGTGGGGT	CGCGTCGAAA	AGCGCGCTTT	AAGGGCGACA	240
	CGCCCCAAAA	GTTTTACCTT	TAAAACTAC	CCGCACGCAG	CACGAACCTG	TTCAGTGATG	300
	TAAATCACCG	CGGAAATATT	GTGGACGTTA	CCCCCGCCTA	CCGCTACGAT	TTCAAAAC	358
20	ATG ACC ATT	TCC TCA CCT	TTG ATT GAC	GTC GCC AAC	CTT CCA GAC	ATC	406
	Met Thr Ile	Ser Ser Pro	Leu Ile Asp	Val Ala Asn	Leu Pro Asp	Ile	
	1	5	10	15			
	AAC ACC ACT	GCC GGC AAG	ATC GCC GAC	CTT AAG GCT	CGC CGC GCG	GAA	454
	Asn Thr Thr	Ala Gly Lys	Ile Ala Asp	Leu Lys Ala	Arg Arg Ala	Glu	
	20	25	30				
25	GCC CAT TTC	CCC ATG GGT	GAA AAG GCA	GTA GAG AAG	GTC CAC GCT	GCT	502
	Ala His Phe	Pro Met Gly	Glu Lys Ala	Val Glu Lys	Val His Ala	Ala	
	35	40	45				
	GGA CGC CTC	ACT GCC CGT	GAG CGC TTG	GAT TAC TTA	CTC GAT GAG	GGC	550
	Gly Arg Leu	Thr Ala Arg	Glu Arg Leu	Asp Tyr Leu	Leu Asp Glu	Gly	
	50	55	60				
30	TCC TTC ATC	GAG ACC GAT	CAG CTG GCT	CGC CAC CGC	ACC ACC GCT	TTC	598
	Ser Phe Ile	Glu Thr Asp	Gln Leu Ala	Arg His Arg	Thr Thr Ala	Phe	
	65	70	75	80			
	GGC CTG GGC	GCT AAG CGT	CCT GCA ACC	GAC GGC ATC	GTG ACC GGC	TGG	646
	Gly Leu Gly	Ala Lys Arg	Pro Ala Thr	Asp Gly Ile	Val Thr Gly	Trp	
	85	90	95				
35	GGC ACC ATT	GAT GGA CGC	GAA GTC TGC	ATC TTC TCG	CAG GAC GGC	ACC	694
	Gly Thr Ile	Asp Gly Arg	Glu Val Cys	Ile Phe Ser	Gln Asp Gly	Thr	
	100	105	110				
	GTA TTC GGT	GGC GCG CTT	GGT GAG GTG	TAC GGC GAA	AAG ATG ATC	AAG	742
	Val Phe Gly	Gly Ala Leu	Gly Glu Val	Tyr Gly Glu	Lys Met Ile	Lys	
	115	120	125				
40	ATC ATG GAG	CTG GCA ATC	GAC ACC GGC	CGC CCA TTG	ATC GGT CTT	TAC	790
	Ile Met Glu	Leu Ala Ile	Asp Thr Gly	Arg Pro Leu	Ile Gly Leu	Tyr	
	130	135	140				
	GAA GGC GCT	GGC GCT CGC	ATT CAG GAC	GGC GCT GTC	TCC CTG GAC	TTC	838
	Glu Gly Ala	Gly Ala Arg	Ile Gln Asp	Gly Ala Val	Ser Leu Asp	Phe	
	145	150	155	160			
45	ATT TCC CAG	ACC TTC TAC	CAA AAC ATT	CAG GCT TCT	GGC GTT ATC	CCA	886
	Ile Ser Gln	Thr Phe Tyr	Gln Asn Ile	Gln Ala Ser	Gly Val Ile	Pro	
	165	170	175				
	CAG ATC TCC	GTC ATC ATG	GGC GCA TGT	GCA GGT GGC	AAC GCT TAC	GGC	934
	Gln Ile Ser	Val Ile Met	Gly Ala Cys	Ala Gly Gly	Asn Ala Tyr	Gly	
	180	185	190				
50	CCA GCC CTG	ACC GAC TTC	GTG GTC ATG	GTG GAC AAG	ACC TCC AAG	ATG	982
	Pro Ala Leu	Thr Asp Phe	Val Val Met	Val Asp Lys	Thr Ser Lys	Met	

EP 0 752 472 A1

		195				200				205							
		TTC	GTT	ACC	GGC	CCA	GAC	GTG	ATC	AAG	ACC	GTC	ACC	GGC	GAG	GAA	ATC
		Phe	Val	Thr	Gly	Pro	Asp	Val	Ile	Lys	Thr	Val	Thr	Gly	Glu	Glu	Ile
5		210						215					220				
		ACC	CAG	GAA	GAG	CTT	GGC	GGA	GCA	ACC	ACC	CAC	ATG	GTG	ACC	GCT	GGC
		Thr	Gln	Glu	Glu	Leu	Gly	Gly	Ala	Thr	Thr	His	Met	Val	Thr	Ala	Gly
		225					230					235				240	
		AAC	TCC	CAC	TAC	ACC	GCT	GCG	ACC	GAT	GAG	GAA	GCA	CTG	GAT	TGG	GTA
10		Asn	Ser	His	Tyr	Thr	Ala	Ala	Thr	Asp	Glu	Glu	Ala	Leu	Asp	Trp	Val
					245						250				255		
		CAG	GAC	CTG	GTG	TCC	TTC	CTC	CCA	TCC	AAC	AAT	CGC	TCT	TAC	ACA	CCA
		Gln	Asp	Leu	Val	Ser	Phe	Leu	Pro	Ser	Asn	Asn	Arg	Ser	Tyr	Thr	Pro
				260					265					270			
		CTG	GAA	GAC	TTC	GAC	GAG	GAA	GAA	GGC	GGC	GTT	GAA	GAA	AAC	ATC	ACC
15		Leu	Glu	Asp	Phe	Asp	Glu	Glu	Glu	Gly	Gly	Val	Glu	Glu	Asn	Ile	Thr
			275						280					285			
		GCT	GAC	GAT	CTG	AAG	CTC	GAC	GAG	ATC	ATC	CCA	GAT	TCC	GCG	ACC	GTT
		Ala	Asp	Asp	Leu	Lys	Leu	Asp	Glu	Ile	Ile	Pro	Asp	Ser	Ala	Thr	Val
		290					295					300					
20		CCT	TAC	GAC	GTC	CGC	GAT	GTC	ATC	GAA	TGC	CTC	ACC	GAC	GAT	GGC	GAA
		Pro	Tyr	Asp	Val	Arg	Asp	Val	Ile	Glu	Cys	Leu	Thr	Asp	Asp	Gly	Glu
		305					310					315				320	
		TAC	CTG	GAA	ATC	CAG	GCA	GAC	CGC	GCA	GAA	AAC	GTT	GTT	ATT	GCA	TTC
		Tyr	Leu	Glu	Ile	Gln	Ala	Asp	Arg	Ala	Glu	Asn	Val	Val	Ile	Ala	Phe
						325					330				335		
25		GGC	CGC	ATC	GAA	GGC	CAG	TCC	GTT	GGA	TTT	GTT	GCC	AAC	CAG	CCA	ACC
		Gly	Arg	Ile	Glu	Gly	Gln	Ser	Val	Gly	Phe	Val	Ala	Asn	Gln	Pro	Thr
			340						345					350			
		CAG	TTC	GCT	GGC	TGC	CTG	GAC	ATC	GAC	TCC	TCT	GAG	AAG	GCA	GCT	CGC
		Gln	Phe	Ala	Gly	Cys	Leu	Asp	Ile	Asp	Ser	Ser	Glu	Lys	Ala	Ala	Arg
			355				360						365				
30		TTC	GTC	CGC	ACC	TGC	GAC	GCG	TTT	AAC	ATC	CCA	ATC	GTC	ATG	CTT	GTC
		Phe	Val	Arg	Thr	Cys	Asp	Ala	Phe	Asn	Ile	Pro	Ile	Val	Met	Leu	Val
			370				375					380					
		GAC	GTC	CCC	GGC	TTC	CTT	CCA	GGC	GCA	GGC	CAG	GAG	TAT	GGT	GGC	ATC
		Asp	Val	Pro	Gly	Phe	Leu	Pro	Gly	Ala	Gly	Gln	Glu	Tyr	Gly	Gly	Ile
			385				390					395				400	
35		CTG	CGT	CGT	GGC	GCA	AAG	CTG	CTC	TAC	GCA	TAC	GGC	GAA	GCA	ACC	GTT
		Leu	Arg	Arg	Gly	Ala	Lys	Leu	Leu	Tyr	Ala	Tyr	Gly	Glu	Ala	Thr	Val
						405					410				415		
		CCA	AAG	ATT	ACC	GTC	ACC	ATG	CGT	AAG	GCT	TAC	GGC	GGA	GCG	TAC	TGC
		Pro	Lys	Ile	Thr	Val	Thr	Met	Arg	Lys	Ala	Tyr	Gly	Gly	Ala	Tyr	Cys
				420					425					430			
40		GTG	ATG	GGT	TCC	AAG	GGC	TTG	GGC	TCT	GAC	ATC	AAC	CTT	GCA	TGG	CCA
		Val	Met	Gly	Ser	Lys	Gly	Leu	Gly	Ser	Asp	Ile	Asn	Leu	Ala	Trp	Pro
			435				440						445				
		ACC	GCA	CAG	ATC	GCC	GTC	ATG	GGC	GCT	GCT	GGC	GCA	GTC	GGA	TTC	ATC
		Thr	Ala	Gln	Ile	Ala	Val	Met	Gly	Ala	Ala	Gly	Ala	Val	Gly	Phe	Ile
			450				455					460					
45		TAC	CGC	AAG	GAG	CTC	ATG	GCA	GCT	GAT	GCC	AAG	GGC	CTC	GAT	ACC	GTA
		Tyr	Arg	Lys	Glu	Leu	Met	Ala	Ala	Asp	Ala	Lys	Gly	Leu	Asp	Thr	Val
			465				470				475				480		
		GCT	CTG	GCT	AAG	TCC	TTC	GAG	CGC	GAG	TAC	GAA	GAC	CAC	ATG	CTC	AAC
		Ala	Leu	Ala	Lys	Ser	Phe	Glu	Arg	Glu	Tyr	Glu	Asp	His	Met	Leu	Asn
					485						490				495		
50		CCG	TAC	CAC	GCT	GCA	GAA	CGT	GGC	CTG	ATC	GAC	GGC	GTG	ATC	CTG	CCA
		Pro	Tyr	His	Ala	Ala	Glu	Arg	Gly	Leu	Ile	Asp	Ala	Val	Ile	Leu	Pro
					500				505					510			

55

EP 0 752 472 A1

AGC GAA ACC CGC GGA CAG ATT TCC CGC AAC CTT CGC CTG CTC AAG CAC 1942  
 Ser Glu Thr Arg Gly Gln Ile Ser Arg Asn Leu Arg Leu Leu Lys His  
 515 520 525  
 5 AAG AAC GTC ACT CGC CCT GCT CGC AAG CAC GGC AAC ATG CCA CTG 1987  
 Lys Asn Val Thr Arg Pro Ala Arg Lys His Gly Asn Met Pro Leu  
 530 535 540  
 TAAATCGGCG AATCCATAAA GGTTCAAAAG AATTCAATAA GGATTCGATA AGGGTTCGAT 2047  
 AAGGGTTCGA TAAGGGCCGA CTTAAATGAT TGGATGTAAA GAAATACCAA TGAAAATTGG 2107  
 10 CAACTCTTTA CACCCAATCT TTAAGACATG GGGGGTGGCG CTGGGCTAAT ATAACCGGTT 2167  
 AGCGAAACGA TTAGTCCCTT GTTAGGGGGA TTAACCCTCG AAGTGGGTCTG TATTTTGGCG 2227  
 TTTGTATGTT CACACAAGAA CCCTGCACAA CGCCTTCAAA GTACGTCGAC CACGACCAAG 2287  
 CGCATATATC ACTCTCACCC TTCAGGATTT AGACTAAGAA ACCATGACTG CAGCACAGAC 2347  
 CAAACCTGAC CTCACCACCA CGGCTGGAAA GCTGTCCGAT CTTGCTCCC GTCTGCAGA 2407  
 AGCTCAAGCT CCAATGGGCG AAGCAACTGT AGAAAAAGTG CACGCTGCTG GCAGGAAGAC 2467  
 15 TGCCCGCGAA CGTATCGAGT ATTTGCTCGA TGAGGGCTCT TTCGTAGAGA TCGATGCTCT 2527  
 TGCTCGTCAC CGTTCCAAGA ACTTCGGCCT GGATGCCAAG CGTCCAGCTA CTGACGGTGT 2587  
 TGTGACTGGT TACGGCACCA TCGATGGCCG TAAGGTCTGT GTGTTCTCCC AGGACGGCGC 2647  
 TGTATTCCGGT GGCCTTTTGG GTGAAGTTTA TGGTGAAGAG ATCGTTAAGG TTATGGATCT 2707  
 TGCGATCAAG ACCGGTGTGC CTTTGATCGG AATCAATGAG GGTGCTGGTG CGCGTATCCA 2767  
 20 GGAAGGTGTT GTGTCTCTGG GTCTGTACTC ACAGATTTTC TACCGCAACA CCCAGGCGTC 2827  
 TGGCGTTATC CCACAGATCT CTTTGATC 2855

INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 543

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: Protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Thr Ile Ser Ser Pro Leu Ile Asp Val Ala Asn Leu Pro Asp Ile  
 1 5 10 15  
 30 Asn Thr Thr Ala Gly Lys Ile Ala Asp Leu Lys Ala Arg Arg Ala Glu  
 20 25 30  
 Ala His Phe Pro Met Gly Glu Lys Ala Val Glu Lys Val His Ala Ala  
 35 35 40 45  
 Gly Arg Leu Thr Ala Arg Glu Arg Leu Asp Tyr Leu Leu Asp Glu Gly  
 50 55 60  
 35 Ser Phe Ile Glu Thr Asp Gln Leu Ala Arg His Arg Thr Thr Ala Phe  
 65 70 75 80  
 Gly Leu Gly Ala Lys Arg Pro Ala Thr Asp Gly Ile Val Thr Gly Trp  
 85 90 95  
 Gly Thr Ile Asp Gly Arg Glu Val Cys Ile Phe Ser Gln Asp Gly Thr  
 100 105 110  
 40 Val Phe Gly Gly Ala Leu Gly Glu Val Tyr Gly Glu Lys Met Ile Lys  
 115 120 125  
 Ile Met Glu Leu Ala Ile Asp Thr Gly Arg Pro Leu Ile Gly Leu Tyr  
 130 135 140  
 45 Glu Gly Ala Gly Ala Arg Ile Gln Asp Gly Ala Val Ser Leu Asp Phe  
 145 150 155 160  
 Ile Ser Gln Thr Phe Tyr Gln Asn Ile Gln Ala Ser Gly Val Ile Pro  
 165 170 175  
 Gln Ile Ser Val Ile Met Gly Ala Cys Ala Gly Gly Asn Ala Tyr Gly  
 180 185 190  
 50 Pro Ala Leu Thr Asp Phe Val Val Met Val Asp Lys Thr Ser Lys Met  
 195 200 205  
 Phe Val Thr Gly Pro Asp Val Ile Lys Thr Val Thr Gly Glu Glu Ile  
 210 215 220

Thr Gln Glu Glu Leu Gly Gly Ala Thr Thr His Met Val Thr Ala Gly  
 225 230 235 240  
 5 Asn Ser His Tyr Thr Ala Ala Thr Asp Glu Glu Ala Leu Asp Trp Val  
 245 250 255  
 Gln Asp Leu Val Ser Phe Leu Pro Ser Asn Asn Arg Ser Tyr Thr Pro  
 260 265 270  
 Leu Glu Asp Phe Asp Glu Glu Glu Gly Gly Val Glu Glu Asn Ile Thr  
 275 280 285  
 10 Ala Asp Asp Leu Lys Leu Asp Glu Ile Ile Pro Asp Ser Ala Thr Val  
 290 295 300  
 Pro Tyr Asp Val Arg Asp Val Ile Glu Cys Leu Thr Asp Asp Gly Glu  
 305 310 315 320  
 Tyr Leu Glu Ile Gln Ala Asp Arg Ala Glu Asn Val Val Ile Ala Phe  
 325 330 335  
 15 Gly Arg Ile Glu Gly Gln Ser Val Gly Phe Val Ala Asn Gln Pro Thr  
 340 345 350  
 Gln Phe Ala Gly Cys Leu Asp Ile Asp Ser Ser Glu Lys Ala Ala Arg  
 355 360 365  
 20 Phe Val Arg Thr Cys Asp Ala Phe Asn Ile Pro Ile Val Met Leu Val  
 370 375 380  
 Asp Val Pro Gly Phe Leu Pro Gly Ala Gly Gln Glu Tyr Gly Gly Ile  
 385 390 395 400  
 Leu Arg Arg Gly Ala Lys Leu Leu Tyr Ala Tyr Gly Glu Ala Thr Val  
 405 410 415  
 25 Pro Lys Ile Thr Val Thr Met Arg Lys Ala Tyr Gly Gly Ala Tyr Cys  
 420 425 430  
 Val Met Gly Ser Lys Gly Leu Gly Ser Asp Ile Asn Leu Ala Trp Pro  
 435 440 445  
 Thr Ala Gln Ile Ala Val Met Gly Ala Ala Gly Ala Val Gly Phe Ile  
 450 455 460  
 30 Tyr Arg Lys Glu Leu Met Ala Ala Asp Ala Lys Gly Leu Asp Thr Val  
 465 470 475 480  
 Ala Leu Ala Lys Ser Phe Glu Arg Glu Tyr Glu Asp His Met Leu Asn  
 485 490 495  
 35 Pro Tyr His Ala Ala Glu Arg Gly Leu Ile Asp Ala Val Ile Leu Pro  
 500 505 510  
 Ser Glu Thr Arg Gly Gln Ile Ser Arg Asn Leu Arg Leu Leu Lys His  
 515 520 525  
 Lys Asn Val Thr Arg Pro Ala Arg Lys His Gly Asn Met Pro Leu  
 530 535 540  
 40

#### 45 Claims

1. A gene derived from a Coryneform bacterium and coding for a protein which imparts surfactant resistance to said bacterium.
- 50 2. The gene according to Claim 1, wherein the protein comprises an amino acid sequence of amino acid number 37 to 543 in an amino acid sequence shown by SEQ ID NO: 2 in the Sequence Listing, or an amino acid sequence having, in said amino acid sequence, substitution, deletion or insertion which does not substantially adversely affect on activity to impart the surfactant resistance to Coryneform bacteria.
- 55 3. The gene according to Claim 1, which comprises a sequence of from 467th to 1987th nucleotides in a nucleotide sequence shown by SEQ ID NO: 1 in the Sequence Listing, or a nucleotide sequence which substantially the same as the sequence.
4. A recombinant DNA obtainable by ligating a vector which functions in Coryneform bacteria to the gene as defined

in Claim 1.

5. A Coryneform bacterium harboring the recombinant DNA as defined in Claim 4.

5 6. A method for producing L-lysine comprising cultivating a Coryneform bacterium which harbors the recombinant DNA as defined in Claim 4 and is capable of producing L-lysine in a liquid medium to produce and accumulate L-lysine in the culture, and collecting the L-lysine.

10 7. A gene comprising a nucleotide sequence having substitution, deletion, insertion, addition or inversion of one or more nucleotides in a nucleotide sequence of the gene as defined in Claim 1 so that a protein encoded by the nucleotide sequence does not normally function regarding an activity to impart surfactant resistance to Coryneform bacteria.

15 8. A recombinant DNA obtainable by ligating a vector which functions in Coryneform bacteria to the gene as defined in Claim 7.

20 9. A Coryneform bacterium having substitution, deletion, insertion, addition or inversion of one or more nucleotides in a nucleotide sequence of the gene as defined in Claim 1 or a promoter thereof on chromosome so that a protein having an activity to impart surfactant resistance to Coryneform bacteria does not normally function.

25 10. A method for producing L-glutamic acid comprising cultivating a Coryneform bacterium having substitution, deletion, insertion, addition or inversion of one or more nucleotides in a nucleotide sequence of the gene as defined in Claim 1 or a promoter thereof on chromosome so that a protein having an activity to impart surfactant resistance to Coryneform bacteria does not normally function, and capable of producing L-glutamic acid, in a liquid medium to produce and accumulate L-glutamic acid in the culture, and collecting the L-glutamic acid.

30

35

40

45

50

55




Fig. 1




Fig. 2




Fig. 3



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP95/00269

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
Int. Cl <sup>6</sup> C12N15/31, C12N1/21, C12P13/08, C12P13/14		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
Int. Cl <sup>6</sup> C12N15/31, C12N1/21, C12P13/08, C12P13/14		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
CAS ONLINE, WPI, WPI/L, BIOSIS PREVIEWS		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP, 5-3793, A (Ajinomoto Co., Inc.), January 14, 1993 (14. 01. 93) & US, 5326693, A & IT, 1244747	1 - 10
A	JP, 53-113086, A (Ajinomoto Co., Inc.), October 3, 1978 (03. 10. 78) (Family: none)	1 - 10
A	JP, 52-24593, B2 (Ajinomoto Co., Inc. and another), July 1, 1977 (01. 07. 77) & FR, 2248320, A1 & US, 3971701, A & GB, 1457953, A & IT, 1022988	1 - 10
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search May 10, 1995 (10. 05. 95)		Date of mailing of the international search report May 30, 1995 (30. 05. 95)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer  Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

THIS PAGE BLANK (USPTO)